







ROUGH CUT

NUGGETS

This Product May Only be Used Inside and Within 100 Feet of Man-Made Structures and For All Burrow Baiting

KILLS RATS, MICE & MEADOW VOLES*



* Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

Net Weight:
4 lbs (1.8 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panels for First Aid and additional precautionary statements.

EPA REG. NO. 12455-81-3240
EPA EST. NOs. 12455-WI-1^K,12455-WI-2^D, 12455-WI-3^W
Superscript is first letter of the lot number.



TO OPEN:

TEAR TOWARD CENTER FROM BOTH NOTCHES

PRESS ZIPPER TO RESEAL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this

IMPORTANT

Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent

- I. Store unused product out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.

 3. Dispose of product container and unused, spoiled, or unconsumed bait as
- specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made structures: House mouse (Mus musculus), Norway pess in and around marimate structures. House mouse (mus musculus), vorway rat (Haftus norvegicus), Roof rat (Haftus raftus), Cotton rat (Sigmodon hispidus), Eastern harvest mouse' (Hafturodontomys humuli), Golden mouse' (Ochrotomys nuttalii), Polynesian rat '(Hattus exulans), Meadow vole' (Microtus pennsylvanicus), White-throated woodrat' (Nectoma albigula), Southern plains woodrat' (Neotoma micropus), and Mexican woodrat' (Neotoma mexicana). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Exemples of such structures include homes and other permanent or infestations. Examples of such structures include homes and other permanent or intestations. Examples or such structures include nomes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, ailerath), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter bailing, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or ford results. food or feed crops.

This product may be applied to active rodent burrows to control rats within or beyond Inis product may be applied to active rodent burrows to control rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active rat burrows. Do not prosecast bait. Because rat infestations may occur in ageas fairther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant balt stations.

Do not sell this product in individual containers holding less than 4 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice or meadow voles' will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as

APPLICATION DIRECTIONS:

RATS: Apply 4 to 16 oz. of bait (usually at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

Мотомсо



KILLS RATS, MICE & MEADOW VOLES

FOR USE IN AND AROUND HOMES AND BUILDINGS

PELLETED BAIT

CTIVE INGREDIENT: Diphacinone (CAS #82-66-6): OTHER INGREDIENTS: 99.995%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this

container. Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

> EPA REG. NO. 12455-81-3240 EPA EST. NOs. 12455-WI-1k, 12455-WI-2P, 12455-WI-3W Superscript is the first letter of the lot number.

Net Weight: 4 lbs (1.8 kg)

Available Exclusively From:



MADE IN USA

The World Leader In Rodent Control Technology®



APPLICATION DIRECTIONS (Continued from other panel)

MICE AND MEADOW VOLES*: Apply 1/4 to 1/2 oz. (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals in infested areas. Larger placements (up to 2 oz.) may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole* activity ease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using waterproof gloves, collect and dispose of all dead, exposed animals and lettover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment, where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and

pels.

All handlers (including applicators) must wear: shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

- · Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
 Do not induce vomiting unless told to do so by the poison control center or doctor.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K_1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-877-854-2494 at once

NOTE TO VETERINARIAN

 $\begin{tabular}{lll} \textbf{Anticoagulant Diphacinone:} & For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K_1. \end{tabular}$

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds and other wildlife. Dogs. cats and other predatory and scavenging mammals and other wingth to poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

080819/9-15



714-1