www.motomco.com



MOTOMOO Medison, WI 53704 U.S.A.



are especially prone to this. These areas will have to be continually monitored and retreated if activity returns. NOTE: At properties are susceptible to recolonization. However, properties near woods, fields, streamstakes, etc. ** Deep tunnel/mound systems require a slightly different procedure. See bottom of package for details.

Five to seven days after application, check for mole activity using the same procedure for determining initial activity. Beturn 49–72 hours later and retreat at active runs.

3 Determine Baiting Results

эспле титикау.

every 5-10 feet of with a small rock. Repeat bait application it not possible, cover pinching hole dosed. Carefully reseal by



the worm should be to his on yewnun arthri the worm is completely opening, ensuring tha Urop one worm into the



larger than the diameter of the WILD & rod slightly Make a note in the





Akik opened runways and revisit them 48-72 hours later. Hunways that have had holes



to not crush runway. subsurface** runways. NOTE: Be careful Use your finger, smal wooden dowel or a narrow rod to puncture a hole in the top of



The Verify Active Runways



Comotolia eeu ot wolf



MOTOMCO FARM & HOME"



MIMICS A MOLE'S NATURAL FOOD SOURCE, THE EARTHWORM



BONUS

1 FREE

Мотомсо **MOLE KILLER**

Other mole baits have been developed on the premise that moles Other mole bails have even developed on the prefines that moles are similar to rats and mice, when in reality moles are insectiones (insect eaters). Moles readily consume earthworms, and when available, grubs. In fact, more than 90% of a mole's average daily dietary requirements are from earthworms and grubs.

Thousands of hours of groundbreaking laboratory and field research have gone into the product you are holding. Motomco Mole Killer has been carefully designed to mimic a mole's most common natural food source – earthworms and grubs. Moles may consume a lethal dose in a single feeding, but it may take two or more days from the time of bait consumption for moles to die.





MOTOMCO

MOLE KILLER

READY-TO-USE

MIMICS A MOLE'S NATURAL FOOD SOURCE, THE EARTHWORM



KEEP OUT OF REACH OF CHILDREN
CAUTION

STORAGE AND DISPOSAL



Мотомсо **MOLE KILLER**

- Scientifically designed and tested to mimic a mole's natural food source. Moles may consume a lethal dose in a single feeding, but it may take two or more days from the time of bait consumption for moles to die.
- Ready To Use
- Effective against most common mole species

FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE	
IF SWALLOWED:	Call a poison control center, doctor, or 1-877-864-2494 immediately for treatment addice. Hare person sip a glass of water if able to swallow. Do not induce vorniting unless told to do so by the poison control center or doctor.
IF ON SKIN:	Wash with plenty of soap and water,



KEEP OUT OF REACH OF CHILDREN CAUTION

See outer container for additional precautionary statements.

FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

IF SWALLOWED:

- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- · Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

IF ON SKIN:

· Wash with plenty of soap and water.

This Product May Only be Used to Control Moles in Manual, Below-Ground Applications.

ACTIVE INGREDIENT:

Bromethalin (CAS #63333-35-7): 0.025%
OTHER INGREDIENTS: 99.975%
TOTAL 100.000%

NET WEIGHT: 10 g

Available Exclusively From:

Мотомсо

3699 Kinsman Blvd., Madison, WI 53704 U.S.A.

www.motomco.com

EPA REG. NO.: 12455-101-3240
EPA EST. NOs. 12455-WI-1k, 12455-WI-2P, 12455-WI-3W
Superscript is first letter of the lot number.

NOTE: MUST BE USED UNDERGROUND

INDIVIDUAL SALE IS PROHIBITED BY LAW

TREATMENT FOR PET POISONING

If an animal eats bait, call veterinarian or 1-877-854-2494 at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the nerve poison Bromethalin. This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

See outer box for full "DIRECTIONS FOR USE" text. See outer box for full "STORAGE AND DISPOSAL" text.