

FIRST AID	
HAVE LABEL WHEN OBTAINING TREATMENT ADVICE	
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person. • Do not induce vomiting unless told to do so by the poison control center or doctor.
NOTE TO PHYSICIAN	
If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K ₁ intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	
TREATMENT FOR PET POISONING	
If an animal eats this bait, call a veterinarian or 1-877-854-2494 at once.	
NOTE TO VETERINARIAN	
Anticoagulant Diphacinone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K ₁ .	

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets.
All handlers (including applicators) must wear shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS
This product is extremely toxic to birds, fish and other wildlife. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

ENDANGERED SPECIES CONSIDERATIONS
NOTICE: It is a Federal offense to use any pesticide product in a manner that results in the death of a member of an endangered species. The use of this product may pose a hazard to Federally designated endangered/threatened species. Suggested measures to protect endangered species in your area may exist in an EPA Interim Measures pamphlet. You may call EPA's Endangered Species Hotline, 1-800-477-3813, to find out if an Interim Measures pamphlet exists in your area and have one sent to you. You can also consult with your local area extension office or pesticide lead agency to determine whether they have imposed any requirements in your area to protect endangered and threatened species.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

Do not sell this product in individual containers holding less than 4 pounds of bait.

USE RESTRICTIONS – For All Uses
Use only in rodent infested areas where exposure to agricultural crops will not occur. Do not contaminate

food or food stuffs. Do not pile bait. Do not apply where contaminating food or surfaces that may contact food is likely. Do not apply bait by aircraft at sites where bait may contaminate bodies of water. When applying bait along canals, and other waterways, use equipment that directs bait away from the water.

NORWAY RATS, ROOF RATS, AND HOUSE MICE

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help prevent exposure:
1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled or unconsumed bait as specified on this label.

USE RESTRICTIONS:
This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Tamper-resistant bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

In active rodent burrows, apply this product to control Norway rats, and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active Norway/roof rat burrows. Do not broadcast bait. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

SELECTION OF TREATMENT AREAS:
Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:
Rats:
Apply 4 to 16 ounces of bait per placement (usually at intervals of 15 to 30 feet). Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

Apply 1 to 2 ounces of bait per placement deep into active rat burrows.

Mice:
Apply ¼ to ½ ounce [1 to 2 level tablespoons] of bait per placement. Space placements at intervals of 8 to 12 feet in infested areas. Larger placements, up to

2 ounces, may be used at points of extremely high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

FOLLOW-UP:
Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all dead, exposed animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish them as needed.

CALIFORNIA GROUND SQUIRRELS

USE RESTRICTIONS:
For control of California Ground Squirrels (*Spermophilus beecheyi*) in industrial, commercial, and residential areas.

Product must be used in a **secured, tamper-resistant bait station** at the following sites: vineyards, fruit tree orchards (stone and pome), nut tree orchards, avocados, and citrus groves; rangeland and pastures; non-crop borders of the listed use sites as follows: outside of fence rows, rights-of-way, canal banks, ditch banks, highways, levees, railroad lines, utilities. It may also be used in and around man-made structures such as agricultural buildings (but not inside of pens), homes, other permanent or temporary structures; campgrounds, golf courses, parks, athletic fields, recreational areas; ornamental plants; lawns; commercial horticultural nurseries, and plantations of forest trees.

Product may also be applied by **spot baiting** applications at the sites listed above, but only during a **non-bearing phase** in vineyards, fruit tree orchards (stone and pome), and nut tree orchards.

Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

BAIT STATIONS:
Place secured tamper-resistant bait stations where California Ground Squirrel activity is present, usually near active burrows. Place bait stations up to 75 feet apart or near active burrows. Apply 2-4 lb. of bait per bait station. Bait stations should be checked every 3-5 days and filled as necessary until signs of bait consumption has ceased.

SPOT BAITING:
Evenly scatter ¼ cup in a measuring cup (0.1 lb.) of bait no larger than 40-50 square feet (i.e. a circular area no more than 8 feet in diameter), around the active California Ground Squirrel burrows and runways. Do not over-bait. Do not place bait in piles. Using the same procedure, make a second application 4 days after the first. Do not apply more than 10 pounds per acre per year.

Surveillance and Follow-Up:
Dead ground squirrels should begin to appear in or near treated areas within 4 to 5 days after the first bait application. Visit the application site to monitor the effects of treatment 4 days after the first application and at subsequent intervals of 1 to 2 days. Wearing gloves, collect and properly dispose of visible carcasses of ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Continue to collect and dispose of dead ground squirrels and search for non-target animals for at least 2 weeks after the last bait application, or longer if carcasses are still being found at that time. Carcass collection and burial should be performed in the late afternoon, near sundown, to reduce the potential for nocturnal, scavengers or predators to find carcasses on the ground surface. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

RESTRICTED USE PESTICIDE
DUE TO HAZARD TO NONTARGET ORGANISMS
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.



KILLS CALIFORNIA GROUND SQUIRRELS, MEADOW & PINE VOLES*, NORWAY RATS, ROOF RATS & HOUSE MICE

**Not approved for the control of Meadow and Pine Voles in the State of California*

ACTIVE INGREDIENT:	
Diphacinone (CAS #82-66-6):	0.005%
OTHER INGREDIENTS:	99.995%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.
[Paper:] Dispose of empty container by placing in trash, or at an approved waste disposal facility or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

See side panels for First Aid, additional Precautionary Statements, and Directions for Use.
See right panels for Health and Safety Sections of this label in Spanish:
Consulte los paneles de la derecha para ver las secciones de Salud y Seguridad de esta etiqueta en español:

NET WEIGHT:
50 lbs (22.68kg)

EPA REG. NO. 12455-136-3240
EPA EST. NOs. 12455-WI-1K, 12455-WI-2P, 12455-WI-3W
Superscript is first letter of the lot number.



Not approved for the control of Meadow and Pine Voles in the State of California
MEADOW AND PINE VOLES

INDUSTRIAL, COMMERCIAL, AND RESIDENTIAL AREAS

USE RESTRICTIONS:
For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*Microtus pinetorum*) in industrial, commercial, and residential areas. Areas include: in and around man-made structures such as agricultural buildings (do not apply in animal pens), homes & lawns (hand baiting into the burrow or tamper-resistant bait station must be used for all residential uses and within 100 feet of a home), other permanent or temporary structures; campgrounds; golf courses; ornamental plants: commercial nurseries. If placed into a tamper-resistant bait station, may also be used in parks, athletic fields, and recreational areas.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

BURROW BAITING:
Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

BAIT STATIONS:
Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

GROUND BROADCAST BAITING:
Broadcast by hand or commercial spreader only within 100 feet of a man-made structure and only in commercial nurseries. Wear waterproof gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later. For pine vole application in states east of the Mississippi River, a rate of 20 lb. per acre may be required. Do not exceed 20 lb. per acre per calendar year.

FRUIT TREE ORCHARDS (DORMANT)

USE RESTRICTIONS:
For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*M. pinetorum*) in pome fruit and stone fruit tree orchards in the **states listed below***. Aerial application is only permitted in the following states where aerial is listed next to the state name. Apply only after fall harvest (including drops), before new spring growth and when no rain or snow is expected for three consecutive days. Do not graze animals in treated areas. ***Connecticut, Georgia, Idaho (aerial), Massachusetts, Michigan (aerial), Missouri, Montana (aerial), North Carolina, New Hampshire, Ohio (aerial), Oregon (aerial), Pennsylvania (aerial), South Carolina, Utah (aerial), Virginia (aerial), Vermont (aerial), Washington (aerial), and West Virginia (aerial).**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

BURROW BAITING:
Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

BAIT STATIONS:
Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

BROADCAST BAITING:
Broadcast by hand, commercial spreader, or aerial (see use restrictions). Wear rubber gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later.

Health and Safety Sections of this label in Spanish:
Secciones de Salud y Seguridad de esta etiqueta en español:

PESTICIDA DE USO RESTRINGIDO
DEBIDO AL PELIGRO PARA ORGANISMOS NO OBJETIVO
Para la venta al por menor y uso únicamente por aplicadores certificados o personas que estén bajo su directa supervisión y únicamente para los usos detallados en el certificado de la Certificación del Aplicador.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN

PRIMEROS AUXILIOS
TENGA LA ETIQUETA CON USTED CUANDO OBTENGA CONSEJOS SOBRE EL TRATAMIENTO

Si se ingiere:

- Llame inmediatamente a un centro de control de envenenamientos, a un médico, o **1-877-854-2494** para consejo de tratamiento.
- Si la persona puede tragar, haga que tome pequeños sorbos de agua.
- No administre nada por vía oral a una persona inconsciente.
- No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico.

NOTA AL MÉDICO
Si se ingiere, este material puede reducir la capacidad de coagulación de la sangre y causar sangrado. En caso de ingestión, administre Vitamina K₁ por vía intramuscular u oral. Repita la administración según sea necesario, según el tiempo de protrombina.

TRATAMIENTO PARA EL ENVENENAMIENTO DE MASCOTAS
Si el animal come el cebo, llame al veterinario o **1-877-854-2494** de inmediato.

NOTA AL VETERINARIO
Anticoagulante Difacinona: Para animales que ingieran cebo y/o muestren signos de envenenamiento (hemorragia o tiempos de protrombina elevados), administre Vitamina K₁.

DECLARACIONES DE PRECAUCIÓN

Riesgos para los seres humanos y animales domésticos
PRECAUCIÓN: Daño si se ingiere. Mantener fuera del alcance de los niños, animales domésticos y mascotas.
Todos los manipuladores (incluidos los aplicadores) deben usar zapatos, calcetines y guantes impermeables. Cualquier persona que recupere cadáveres o cebo no utilizado después de la aplicación de este producto debe usar guantes impermeables.

Requerimientos de seguridad para el usuario
Siga las instrucciones del fabricante para la limpieza/mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separadamente de otra ropa para lavar. Los usuarios deben quitarse inmediatamente el equipo de protección personal (PPE) luego de manipular este producto. Lave el exterior de los guantes antes de quitárselos. Tan pronto sea posible, Lávese con abundante agua y jabón luego de manipular los pesticidas y antes de comer, beber, masticar chicle, usar tabaco, o ir al baño.

PELIGROS AMBIENTALES
Este producto es extremadamente tóxico para aves, peces y otros animales salvajes. Perros, gatos y otros mamíferos y aves depredadores y carroñeras podrían intoxicarse si se alimentan de animales que han ingerido este cebo. No aplique directamente a cuerpos de agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. No contamine cuerpos de agua cuando disponga de las aguas de lavado de equipo aguas de enjuague.

INSTRUCCIONES DE USO
Es una violación de la ley federal utilizar este producto de manera inconsistente con lo indicado en su etiqueta.

LEA ESTA ETIQUETA: Lea esta etiqueta completa y siga todas las instrucciones y precauciones de uso. Use únicamente para los sitios, plagas y métodos de aplicación descritos en esta etiqueta.
No venda este producto en envases individuales que contengan menos de 4 libras de cebo.

ALMACENAMIENTO Y DESECHO
No contamine agua, comida, ni forrajes mediante el almacenamiento o desecho.
ALMACENAMIENTO DE PESTICIDAS: Almacene solo en el envase o recipiente original, en un lugar fresco y seco, fuera del alcance de niños y mascotas. Mantenga los envases cerrados y alejados de otros productos químicos.
DESECHO DE PESTICIDAS: Deseche los desechos resultantes del uso de este producto en la basura o en una instalación de eliminación de desechos aprobada.
MANIPULACIÓN DE RECIPIENTES: Recipiente no rellenable. No reutilice o rellene este recipiente.
[Papel:] Deseche el envase vacío en la basura, en un punto de recogida de residuos autorizado o incinerándolo. En la mayoría de los estados, no se permite la quema.

